Mon. Not. R. Astron. Soc. **000**, 1–14 (2010) Printed 21 January 2013 (MN LATEX style file v2.2)

The Padova-Millennium Galaxy and Group Catalogue (PM2GC): the group-finding method and the PM2GC catalogues of group, binary and single field galaxies

Rosa Calvi[1⋆], Bianca M. Poggianti[2], Benedetta Vulcani[1,2]
[1]*Astronomical Department, Padova University, Italy*
[2]*INAF-Astronomical Observatory of Padova, Italy*

arXiv:1105.3683v1 [astro-ph.CO] 18 May 2011

Accepted Received ..

ABSTRACT

We present the construction and describe the properties of the Padova-Millennium Galaxy and Group Catalogue (PM2GC), a galaxy catalogue representative of the general field population in the local Universe. We characterize galaxy environments by identifying galaxy groups at $0.04 \leqslant z \leqslant 0.1$ with a Friends-of-Friends (FoF) algorithm using a complete sample of 3210 galaxies brighter than $M_B = -18.7$ taken from the Millennium Galaxy Catalogue (MGC, Liske et al. (2003)), a $38 \mathrm{deg}^2$ photometric and spectroscopic equatorial survey. We identified 176 groups with at least three members, comprising in total 1057 galaxies and representing $\sim$43 per cent of the general field population in that redshift range. The median redshift and velocity dispersion of our groups are 0.0823 and 192 $\mathrm{km\,s^{-1}}$, respectively. 88 per cent of the groups have fewer than ten members, and 63 per cent have fewer than five members. Non-group galaxies were subdivided into "binary" systems of two bright close companions, and "single" galaxies with no companion, in order to identify different environments useful for future scientific analysis. We performed a detailed comparison with the 2PIGG catalog to validate the effectiveness of our method and the robustness of our results. Galaxy stellar masses are computed for all PM2GC galaxies, and found to be in good agreement with Sloan Digital Survey Data Release 7 (SDSS-DR7) mass estimates. The catalogues of PM2GC groups, group properties and galaxy properties in all environments are publicly available on the World Wide Web.

Key words: galaxies: formation: general – galaxies: groups environments – galaxies: stellar masses

1 INTRODUCTION

In the ΛCDM model of the Universe, the growth of large-scale structure occurs hierarchically with the most massive structures, like clusters and superclusters, formed through the accretion of smaller halos which continuously interact and merge, acquiring infalling galaxies along filaments.

Historically, the major discoveries concerning the impact of hierarchical assembly and thus environment on galaxy evolution come from the studies conducted on clusters. The Butcher-Oemler effect, the morphology vs. density relation and the star formation vs. density relation have shown that the properties of galaxies within clusters are strongly correlated with the local environment and evolve with redshift (see, e.g., Dressler 1980; Butcher & Oemler 1984; Dressler et al. 1997; Poggianti et al. 1999; Balogh & Bower 2003). In agreement with the observed trends a presumable scenario is that the clustering process itself would drive the evolution of the galaxy properties and the typical increase of the early-type fraction with decreasing redshift would be due to gas-rich, star-forming disk galaxies which fall into clusters at higher redshift (z$\sim$0.5-1) and have their gas reservoir depleted by some mechanism that transforms them into red, bulge-dominated and quiescent galaxies at z$\sim$0.

Among the numerous mechanisms proposed to deplete the reservoir of gas in late-type galaxies are ram-pressure and/or tidal stripping (Gunn & Gott 1972), interaction with the cluster potential (Byrd & Valtonen 1984) and repeated high-velocity encounters ("harassment") (Moore et al. 1996). However, recent results (Zabludoff & Mulchaey 1998; Mulchaey et al. 2003) have focussed on the hypothesis that the anomalous fraction of early-type galaxies in clusters is the consequence of some pre-process which takes place in *groups* before galaxies are accreted into the cluster. This hypothesis is supported by the observations of a decline in star formation rate (SFR) in the outskirts of clusters, well outside the virial radius (Balogh et al. 1999; Lewis et al. 2002). Although

⋆ E-mail: rosa.calvi@unipd.it

Berrier et al. (2009) found that the pre-processes in the group environment seem not to produce a large effect, it is nevertheless true that the mechanisms like galaxy mergers and strangulation that efficiently act also in groups could play a very important role in the formation of the galaxy populations, as it has been shown by combining high-resolution N-body simulations with semi-analytic models for galaxy evolution (Springel et al. 2001; Kang et al. 2005; Mihos 2004; Toomre, A. & Toomre, J. 1972; Barnes & Hernquist 1996; Cox et al. 2008; Murante et al. 2007; Somerville et al. 2008; Guo et al. 2011; Wang, Kauffman & De Lucia 2007; Font et al. 2008; Weinmann et al. 2010).

Compared to clusters galaxy groups are more difficult to detect because they have a lower density with respect to the background galaxy population, but are much more common in the Universe. Today over 50 per cent of galaxies are in these systems and span a wide range in local density showing properties which range from cluster-like to field-like (Zabludoff & Mulchaey 1998). For this reason they are thought to be a key environment to investigate galaxy evolution and to provide a clear framework on the nature of the physical galaxy transformation mechanisms.

Until recently, the difficulties in obtaining large, unbiased samples of groups have forced most of the studies to use small samples selected, for example, from the Hickson compact group catalogue (Hickson, Kindl & Auman 1989), from the CfA redshift survey (Geller & Huchra 1983; Moore, Frenk & White 1993), and from X-ray surveys (Henry et al. 1995; Mulchaey et al. 2003). Only with the advent of large galaxy redshift surveys, such as the Two degree Field Galaxy Redshift Survey (2dFGRS), the Sloan Digital Sky Survey (SDSS) and the Canadian Network for Observational Cosmology Redshift Survey (CNOC2), has it become possible to generate large group catalogues in the local Universe (e.g. Huchra & Geller 1982; Ramella, Geller & Huchra 1989; Ramella, Pisani & Geller 1997; Ramella et al. 1999; Hashimoto et al. 1998; Tucker et al. 2000; Martínez et al. 2002; Eke et al. 2004; Balogh et al. 2004), and at intermediate redshift (Carlberg et al. 2001; Wilman et al. 2005).

The methods used to identify galaxy groups depend on several assumptions and there is no unique group-finder algorithm to assign them members. The most popular is the Friends-of-Friends (FoF) algorithm applied to galaxy redshift surveys, first proposed by Huchra & Geller (1982). Other approaches, which strongly depend on the properties of dark matter haloes, can be found in Marinoni et al. (2002); Gerke et al. (2005); Miller et al. (2005); Yang et al. (2005). The 2PIGG (2dFGRS Percolation-Inferred Galaxy Group) catalogue (Eke et al. 2004) and the SDSS group catalogue (Berlind et al. 2006) are two of the largest available samples of galaxy groups which use "realistic" mock catalogues to calibrate the parameters associated with the group-finder algorithm.

The motivation for another group catalogue and for our work is to provide a new dataset characterized by both high spectroscopic completeness, to define galaxy environment well, with high quality imaging, to investigate galaxy properties such as galaxy morphologies which could not explored in detail in other catalogues.

In this paper we present the construction and describe the properties of the Padova-Millennium Galaxy and Group Catalogue (PM2GC), a database of groups and galaxies at low redshift fully provided, easily upgradable and easily to be consulted. The PM2GC redshift range ($0.03 \leqslant z \leqslant 0.11$) is similar to WINGS (WIde-field Nearby Galaxy-cluster Survey), a survey of 77 X-ray selected galaxy clusters (Fasano et al. 2006). The combination of PM2GC and WINGS allows the study of the properties and the evolution of galaxies in the widest possible range of environments in the local Universe and to understand the origin of the observed trends of galaxy properties versus environment and the relation between galaxy star formation histories and the growth history of structures.

This paper is structured as follows. In §2 we present the dataset (MGCz) used and give an overview of the survey; in §3 we provide a description of our approach to identify groups and our group-finder algorithm linking criteria; in §4 we show the properties of groups and galaxies in the different environments identified; in §5 we derive the galaxy stellar masses comparing them with SDSS-DR7 masses; in §6 we test the reliability of our group sample comparing it with the 2dFGRS group catalogue (2PIGG, Eke et al. 2004); in §7 we present the publicly released PM2GC catalogues and finally in §8 we present a summary. Throughout the paper we adopt $H_0 = 70\,\mathrm{km\,Mpc^{-1}s^{-1}}$, $h = H_0/100$, Ω_m =0.3 and Ω_λ =0.7. All magnitudes are expressed in the Vega system unless otherwise stated.

2 THE DATA

To build a catalogue that satisfies our requirements of spectroscopic and photometric completeness, we used a set of galaxies derived from the Millennium Galaxy Catalogue (MGC) (Liske et al. 2003; Cross et al. 2004; Driver et al. 2005), a B-band imaging survey, both deep and wide, which provides a high quality, complete representation of the nearby galaxy populations.

A detailed description of the survey strategy, the photometric and astrometric calibration and the object detection and classification can be found in Liske et al. (2003). In brief, the survey extends along an equatorial strip covering an area of $\sim$37.5 deg^2 and consists of 144 overlapping fields taken with the WFC four-CCD mosaic on the Isaac Newton Telescope, with a uniform isophotal detection limit of 26.0 mag arcsec^{-2}. The catalogue contains about one million of objects reduced by the Cambridge Astronomy Survey Unit (CASU) (Irwin & Lewis 2001) and classified using Source Extractor (SExtractor, Bertin & Arnouts (1996)). The entire set of objects, spanning the range $16 \leqslant B_{MGC} < 24$, was next divided into two magnitude ranges to better address the division between stars and galaxies: the MGC-BRIGHT catalogue, which contains all objects with $B_{MGC} < 20$ mag, and the MGC-FAINT catalogue which contains the others.

For this paper we selected a sample of galaxies from the MGCz catalogue - a version of the total MGC database available on DVD - that is the spectroscopic extension of MGC-BRIGHT. It was built upon the redshifts provided by the 2dFGRS and the SDSS-EDR/DR1, in which the MGC survey region is fully contained, and completed with redshifts taken by the MGC team at the Anglo Australian Telescope using the 2dF facility (Driver et al. 2005), as well as redshifts from the NASA Extragalactic Database (NED), the 2dF QSO Redshift Survey (2QZ), the Paul Francis' Quasar Survey and of some low surface brightness galaxies. The total spectroscopic completeness of galaxies obtained by MGC team is greater than 96 per cent for $B_{MGC} < 20$, so we have no need to apply a statistical completeness correction to sample[1].

At the beginning, we extracted galaxies at $0.03 \leqslant z \leqslant 0.11$; we chose this redshift range to avoid galaxies too close by whose spectra only sample the central regions, while remaining at sufficiently

[1] For the additional completeness test see §3

Environment	Number of galaxies	
	$0.04 \leqslant z \leqslant 0.1$	$0.03 \leqslant z \leqslant 0.11$
Groups	1057	–
field-single	846	1141
field-binary	367	490
Mix sample	208	522
General field	2460	3210

Table 1. List of the number of galaxies in the different environments.




Figure 1. Absolute magnitude in B band vs. redshift for our data sample.

low redshifts to retain a deep absolute magnitude completeness limit. Absolute B-band magnitudes were obtained k-correcting the observed SExtractor 'BEST' magnitudes (MAGAUTO, except in crowded region where the ISOCOR magnitude was used instead), corrected for Galactic extinction. The k-corrections were taken from Poggianti (1997), using the galaxy redshift and the Sloan galaxy color provided in the MGCz catalogue (hereafter, MGC-SDSS).

To build our catalogues we used only 3210 bright galaxies with a magnitude M_B <-18.7 corresponding to the k-corrected B_{MGC} =20 magnitude at our redshift upper limit. In Fig.1 a plot of absolute magnitude vs. redshift for this sample is shown. The high spectroscopic completeness to B_{MGC} <20, coupled with the photometric depth, makes it a complete absolute magnitude-limited sample.

3 GROUP BUILDING METHOD

The approach we used to identify galaxy groups is similar to that adopted by McGee et al. (2008), and is based on a plain FoF algorithm. According to FoF criteria, two galaxies, *i* and *j*, are physically related and join the same group if their distances in the projected direction (D) and in the line-of-sight (V) are less than some fixed thresholds, i.e.

$$D_{ij} \leqslant D_L; \quad V_{ij} \leqslant V_L \tag{1}$$

D_L and V_L are called "linking lengths" and link together all galaxies within a particular linking volume. We chose these lengths to take into account the typical gravitational bounds of groups and to follow a similar approach of that used to identify groups at high redshift in the ESO Distant Cluster Survey (White et al. 2005; Halliday et al. 2004; Milvang-Jensen et al. 2008; Poggianti et al. 2006). Similarly to other studies (McGee et al. 2008; Wilman et al. 2005), we adopted for a linking volume a cylinder centred on each galaxy with radius

$$D_L = 0.5 h^{-1} Mpc \tag{2}$$

corresponding to a density contour $\frac{3}{4\pi D_L^3 n} = 216.6 gal^{-1}$, being n the mean observed number density of galaxies in the total sample. The line-of-sight depth V_L is equal to three times the velocity dispersion, fixed at 500 $\mathrm{km\,s^{-1}}$ rest frame, of the galaxy redshift.

For each galaxy in our sample brighter than M_B =-18.7, we obtained its first neighbours in the cylinder defined above, and added to these, by a recursive procedure, neighbors of neighbors, until no more are found. The resulting system we defined the "trial" group. Only systems with at least three galaxies were further considered as group candidates.

As a second step, we computed for each trial group its median geometric centre, median redshift and velocity dispersion using the statistical methods by Beers, Flynn & Gebhardt (1990), considering the gapper scale estimator for groups with less than ten galaxies and the biweighted scale estimator for more populous systems. A galaxy was considered member of a group if its spectroscopic redshift lay within $\pm 3\sigma$ from the median group redshift and if it was located within a projected distance of $\pm 1.5\, R_{200}$ from the geometrical centre. R_{200} is an approximation of the virial radius - the radius which delimites a sphere with mean interior density 200 times the critical density - computed as in Finn et al. (2005)

$$R_{200} = \frac{1.73\sigma}{1000 kms^{-1}\sqrt{\Omega_\Lambda + \Omega_0 (1+z)^3}} h^{-1} Mpc \tag{3}$$

where σ and z are the group's velocity dispersion and median redshift, respectively.

We iterated the second step several times, recalculating every time the group redshift, velocity dispersion and R_{200}, and moving to the next iteration only those groups with at least three members. The process stops when the last two iterations have identical output. At most, three iterations were sufficient to reach convergence. We consider members of the final groups only those galaxies that are within $1.5\, R_{200}$ from the group centre and 3σ from the group redshift.

Using this method we obtained a sample of 176 galaxy groups in the redshift range 0.04⩽z⩽0.1 containing in total 1057 group members with magnitude M_B <-18.7.

Groups below z∼0.04 and above z∼0.1 are disregarded in the following analysis, because, for a maximum velocity dispersion of 800 $\mathrm{km\,s^{-1}}$, and due to the redshift limits of our original sample $(0.03 - 0.11)$, they can suffer from spectroscopic incompleteness. We have mentioned before that the total spectroscopic completeness of our sample is 96% to $B = 20$. Sky regions with the highest galaxy density, such as groups, might in principle suffer from higher incompleteness, due to the difficulty to place fibers close together. This problem is strongly mitigated in the MGC sample, because it is a combination of three different spectroscopic campaigns (SDSS, 2dF and MGCz). However, in order to double-check the completeness in groups, we performed an additional test. For each PM2GC group, within the angular radius corresponding to R_{200}




Figure 2. Redshift (z) distribution (left), velocity dispersion (σ) distribution (centre) and number of members distribution (right) of the 176 groups at $0.04 \leqslant z \leqslant 0.1$.

from the group's centre, we counted the number of MGC galaxies with redshift and the total number of galaxies in the photometric catalog brighter than $B = 20$. The ratio of the sum of all galaxies with redshift and the sum of all galaxies in the photometric sample for all groups is r_1. Using the same R_{200} we considered for each group a random RA, DEC for the group centre within the MGC area and counted again the number of galaxies with redshift and the number in the photometric catalog brighter than $B = 20$ within the same R_{200}. For each group we repeated this 100 times, finding each time a value r_2. The mean value of the ratio r_1/r_2 is 0.999 confirming the high spectroscopic completeness of the galaxy redshift catalogue even for galaxies preferentially clustered in groups.

Moreover, we analyzed our 176 groups to assess whether they are fully contained in the narrow strip of the MGC survey: the aim was to understand how many and which groups suffer from edge problems and therefore need to be treated with caution in the subsequent analysis. Looking at the group centre position coordinates, we flagged those 66 groups for which $RA_{centre} \pm R_{200}$ and/or $DEC_{centre} \pm R_{200}$ fall out of the ranges of the MGC strip.[2] Six of these had clear edge problems also from the comparison with 2PIGG (see §6).




Figure 3. Redshift distribution for the different samples of galaxies: 1141 field-single galaxies (dotted line, $0.03 \leqslant z \leqslant 0.11$), 490 field-binary galaxies (dot-short dashed line, $0.03 \leqslant z \leqslant 0.11$), 1057 group galaxies (solid thin line, $0.04 \leqslant z \leqslant 0.1$) and 3210 general field (GF) galaxies (solid thick line, $0.03 \leqslant z \leqslant 0.11$).

4 OTHER ENVIRONMENTS

Galaxies that are not members of our 176 groups are treated separately, to study galaxy properties in several environments and compare the results.

We named "field-single" and "field-binary" those subsets of galaxies at $0.03 \leqslant z \leqslant 0.11$ that have no friends (1141) or solely one friend (490) in their original trial cylindrical volume, respectively. The first sample, which contains isolated galaxies, is considered as pure field; the second one is composed of binary systems of galaxies, i.e. those pairs of bright galaxies that have a projected mutual separation within $0.5h^{-1}$Mpc and a redshift within $1500\mathrm{km\,s^{-1}}$.

The remaining 522 galaxies that are not in groups, field-single or field-binary environments are either those that are in groups at $z<0.04$ or $z>0.1$ (302) or those galaxies that, although located in a trial group, did not make it into the final group sample (220). These galaxies are part of the outer regions of groups (outside $1.5\,R_{200}$), therefore we prefer not to consider them as "single".

Finally, the sample of all galaxies in all environments at $0.03 \leqslant z \leqslant 0.11$ is named "general field" (GF from now on), and is representative of the general field low-z galaxy population. In Table 1 we list the number of galaxies in different environments.

5 PROPERTIES OF GROUPS AND GALAXIES IN THE DIFFERENT ENVIRONMENTS

In Fig.2 we show the general characteristics of our group sample. It is clear that, as in other catalogues, most of the groups lie in the higher redshift range, and contain fewer than 10 members.

[2] The fractional area lost adopting these limits instead of $RA_{centre} \pm 1.5\,R_{200}$ and $DEC_{centre} \pm 1.5\,R_{200}$ is negligible, therefore also the difference in number of galaxies falling outside of the field is irrelevant.




Figure 4. Left. B-band absolute magnitude distribution for the different samples of galaxies: 1141 field-single galaxies (dotted line, 0.03⩽z⩽0.11), 490 field-binary galaxies (dot-short dashed line, 0.03⩽z⩽0.11), 1057 group galaxies (solid thin line, 0.04⩽z⩽0.1) and 3210 general field (GF) galaxies (solid thick line, 0.03⩽z⩽0.11). **Right.** The absolute magnitude distributions of group, binary and field galaxies, all in the range 0.04⩽z⩽0.1, normalized to the same total number of galaxies (N=1000).

The median redshift and velocity dispersion of the sample are 0.0823 and 191.8 $\mathrm{km\,s^{-1}}$, respectively. The range of velocity dispersion is between 100 $\mathrm{km\,s^{-1}}$ and 800 $\mathrm{km\,s^{-1}}$for most groups, with 11 per cent having a velocity dispersion $<$100 $\mathrm{km\,s^{-1}}$ and 29 per cent $>$400$\mathrm{km\,s^{-1}}$. Hence, a significant fraction of the structures we identify have velocity dispersions higher than 400 $\mathrm{km\,s^{-1}}$, which is the commonly adopted limit between groups and clusters. The fraction of groups with less than five members is 63 per cent and 43 per cent have only three members.

In Fig.3 we show a comparison of the redshift distribution of galaxies in the several environments we have identified. We note the presence of a prominent peak at z∼0.095 in the general field distribution, due to groups likely belonging to a quite populated structure at that redshift.

The magnitude distribution of galaxies in the different environments is shown in Fig.4. Raw numbers are given in the left panel, while in the right panel the group, binary and single galaxy distributions, all in the range 0.04⩽z⩽0.1, have been normalized to the same number of galaxies (N=1000) to show the differences. From this figure, the relative proportion of faint galaxies in the single and binary fields seems higher than in groups.




Figure 5. Galaxy stellar mass distribution for the different samples of galaxies: field-single (dot line, 0.03⩽z⩽0.11), field-binary (dot-short dash line, 0.03⩽z⩽0.11), groups (solid thin line, 0.04⩽z⩽0.1) and general field (GF) (solid thick line, 0.03⩽z⩽0.11) galaxies.

6 GALAXY STELLAR MASSES

We determined the stellar masses for all galaxies in our sample using the Bell & de Jong (2001) relation according to which, under the assumption of a universal IMF, the stellar mass-to-light (M/L) ratio is strongly correlated with the optical colors of the integrated stellar populations. Using the B-band photometry, taken from MGCz, we apply the equation

$$\log_{10}(M_\star/L_B) = a_B + b_B(B-V) \qquad (4)$$




Figure 6. Comparison between the masses of galaxies in our total sample determined, in this paper, using the Bell & de Jong (2001) relation and g and r MGC-SDSS magnitudes, and the masses of the same galaxies found in the SDSS-DR7 catalog. The black line is the 1:1 relation.

having considered a Bruzual & Charlot model with a_B=-0.51 and b_B=1.45 for a Salpeter (1955) IMF (0.1-125 M$_\odot$) and solar metallicity. To compute the rest frame $(B-V)$ color we followed the filter conversions from Blanton & Roweis (2007), i.e.

$$(B-V)_{AB} = 0.5928 + 1.1521[(g-r) - 0.6148] \quad (5)$$

using the MGC-SDSS (from SDSS-EDR/DR1) $(g-r)$ color corrected for extinction. We then added 0.11 to the $(B-V)$ colors to transform them from the AB system to the Vega system, and applied the k-corrections in B and V to obtain the rest frame $(B-V)$ colors. The galaxy stellar masses found with the eq.(4) were subsequently scaled to a Kroupa (2001) IMF to compare with the SDSS, using a conversion factor from Salpeter to Kroupa of 1/1.55 (Cimatti et al. 2008). We also took into account the fact that a certain number of galaxies lie in regions where the photometry can be affected by CCD edges, satellite trails, bright stars and galaxies (B$_{MGC}$ <12.5), diffraction spikes and so on; any object in these regions was marked by a flag in the MGCz catalogue to indicate that it may have an incorrect photometry. Comparing the B$_{MGC}$ magnitude with the B$_{MGC-SDSS}$ magnitude (determined using the MGC-SDSS color $g-r$), we have used the B$_{MGC-SDSS}$ magnitude to determine the stellar masses for those galaxies for which ΔB = |B$_{MGC}$-B$_{MGC-SDSS}$ | > 0.5 mag.

The histogram of the mass distribution for galaxies brighter than M_B =-18.7 in the different environments is shown in Fig.5. As discussed in Calvi et al. (2011b) (in preparation), the mass completeness limit for our sample is $M = 10^{10.25} M_\odot$, so any meaningful comparison must be done above this limit. The variation of the mass function with environment will be discussed in Calvi et al. (2011b).

We also compared our mass estimates with the stellar masses computed from the Sloan collaboration from the SDSS-DR7 catalogue[3] for those galaxies whose MGC and DR7 positions match within 1 arcsec. DR7 masses are computed based on the Sloan photometry, using model magnitudes, for a Kroupa IMF in the range 0.1-100 $M_\odot$ (J. Brinchmann 2010, private communication).

In Fig.6 we show this comparison for the GF galaxies in common with the DR7 mass catalogue. The agreement is satisfactory at masses above log$_{10}$(M$_\star$/M$_\odot$) ∼10.3. Here, the dispersion is similar to the typical mass error for our method that is normally taken to be 0.2-0.3 dex. At lower masses, there is a systematic effect, surely due to the different mass estimate methods, in the sense that our masses are higher than SDSS-DR7 masses by up to $\sim 0.2-0.3$ dex.

Finally, with the Bell & de Jong (2001) method, we also computed the stellar masses using the SDSS DR7 model magnitudes corrected for extinction, obtained from the SDSS Catalogue Archive Server (CAS) system for 3140 of our galaxies, and verified that these masses were in very good agreement with those based on MGC-SDSS magnitudes, without any offset at low masses (plot not shown).

7 DATASET VALIDATION

The construction of a robust catalogue of groups is essential to characterise accurately their properties. To validate our catalogue, we concentrated on a direct comparison with one of the largest galaxy group sample, the 2PIGG catalogue (Eke et al. 2004). This consists of ∼290000 groups, with at least two members, found in the Northern and Southern Galactic Patches (NGP and SGP) in the 2dFGRS using a group finding procedure based on a FoF algorithm. They found galaxy groups using linking parameters calibrated on realistic mock catalogues identified with high-resolution N-body simulations and a semi-analytical model of galaxy formation. Their purpose was to provide overdensity regions that have velocity dispersions and projected sizes similar to those of their parent dark matter halos.

The comparison with the 2PIGG catalogue is an important step to test the work assumptions we made in the FoF algorithm and validate the effectiveness of our sample. Our galaxy groups are all contained in their Northern Galactic Patch.

To match our groups with the 2PIGG catalogue, we checked when:

(i) the geometric centres agreed within 0.1°;
(ii) the group redshifts differed by at most 0.0007.

81 of our groups satisfy the above criteria and match a 2PIGG group. For these, in Fig.7 we show the comparison between our and 2PIGG redshifts (left panel) and velocity dispersions (right panel). The 2PIGG velocity dispersions used in this plot are gapper estimates, derived from their tabulated σ values using eq.(4.6) in Eke et al. (2004). 2PIGG adopted a fixed error on σ of 85 $\mathrm{km\,s^{-1}}$, which is displayed in Fig.7. For some 2PIGG groups the sigma value is 0, which means that the individual galaxy error is at least as big as their estimate of the velocity dispersion. As expected, given the large uncertainty in the σ measurements based on a few redshifts, there is a large scatter in this comparison, although 75 per cent are within the errors.

As a further step, we performed a match between the position of our group galaxies and that of 2PIGG galaxies within 3 arcsec. All of these matches agreed also in redshift. This allowed us

[3] http://www.mpa-garching.mpg.de/SDSS/DR7/Data/stellarmass.html

to investigate if our group galaxies were associated with the same 2PIGG group matching in geometric centre. In this way we also found how many of our group galaxies were observed by 2PIGG.

For 23 of our groups none or fewer than 50 per cent of our galaxies have been observed by 2PIGG. Not surprisingly, these groups do not match any 2PIGG group according to criteria (1) and (2) above.

For another 20 of our groups we did not find a match in both barycentre and redshift with 2PIGG even if at least 50 per cent of our galaxies have been observed by 2PIGG. However, we have a higher number of redshifts than 2PIGG in most of these groups.

The remaining 52 of our 176 groups have peculiar characteristics and deviate from 2PIGG groups. 34 were associated with one group for 2PIGG but for us they were split in two or more groups; 11 groups have a high velocity dispersion for their number of galaxies; for the remaining 7 groups, the barycentre of the corresponding 2PIGG group falls out of the MGC survey strip, showing that these groups are affected by edge problems in the MGC thus they will be disregarded in our analysis. To conclude, about half of the PM2GC groups have a correspondence in the 2PIGG catalogue. In addition, we found a number of groups that 2PIGG did not identify. This is due to the higher spectroscopic completeness of the PM2GC, that contains 1074 galaxies (33 per cent of the PM2GC catalogue) that were not observed by the 2dFGRS. Moreover, only 20 per cent of our field-single galaxies belong to a group according to 2PIGG, confirming the overall statistical agreement with 2PIGG.

8 PM2GC PRESENTATION

In this last section we describe the five main catalogues we provide: the group catalogue, and the catalogues of galaxies in groups, in binary systems, in single field environment and in the general field.

We stress that the group catalogue contains only groups whose redshift lies at $0.04 \leqslant z \leqslant 0.1$ because, given our selected galaxy redshift range $0.03 \leqslant z \leqslant 0.11$ and for a maximum velocity dispersion of about 800 $\mathrm{km\,s^{-1}}$, groups at z below $\sim$0.04 and above $\sim$0.1 could be incomplete. In contrast, the general field, field and binary catalogues contain all galaxies in the range $0.03 \leqslant z \leqslant 0.11$.

The main properties of a subsample of galaxy groups in the PM2GC catalogue are given in Table 2 (corrisponding to Table 2 in electronic version for the total sample). The different columns indicate: (1) PM2GC group serial number; (2) median group redshift; (3) rest frame velocity dispersion ($\mathrm{km\,s^{-1}}$) computed as in §3; (4) velocity dispersion error ($\mathrm{km\,s^{-1}}$); (5) number of galaxies contained within a projected radius $1.5\,R_{200}$ from the group geometrical centre and within 3σ from the group redshift; (6-7) geometric centre right ascension and declination at epoch J2000 (in degrees); (8) R_{200} in Mpc, (9) matched 2PIGG group serial number (9999 if no match with 2PIGG); (10) 2PIGG group, or groups, to which galaxies of our group belong; (11) number of galaxies of our group that are members of the 2PIGG matched group (0 indicate no galaxies in 2PIGG or ungrouped galaxies); (12) 2PIGG match flag (1=good match with 2PIGG, both in geometric centre and $\Delta z \leqslant 0.0007$, one to one group correspondance, all of our galaxies in common with 2PIGG are in the matched 2PIGG group; 2= as 1 but at least one of our group galaxies is either ungrouped for 2PIGG, and/or belongs to a different 2PIGG group, and/or is not present in 2PIGG; 3 = no match with 2PIGG, 2PIGG has none or < 50 per cent of our redshifts; 4= no match with 2PIGG, and 2PIGG has at least 50 per cent of our redshifts; 5= one group for 2PIGG, two or more groups for us; 6= no match with 2PIGG, high σ for number of members; 0= group with likely edge problems by comparison with 2PIGG), (13) Edge flag, see §3 1=no edge problem; 2= edge problem.

In the galaxy catalogues we list the properties of galaxies in different environments. In the catalogue of galaxies in groups, see Table 3 for a subsample of galaxies (corrisponding to Table 3 in electronic version for the total catalogue), the columns indicate: (1) galaxy serial number in MGC; (2) galaxy redshift in MGC; (3-4) right ascension and declination in MGC at epoch J2000 (in degrees); (5) group number in PM2GC; (6) distance from group geometric centre in Mpc; (7) distance from group geometric centre in R_{200} units; (8) logarithm of galaxy stellar mass (Kroupa IMF) obtained from MGC-SDSS colors (PM2GC mass); (9) logarithm of galaxy stellar mass (Kroupa IMF) we computed using DR7-CAS colors; (10) logarithm of galaxy stellar mass (Kroupa IMF) given in the SDSS-DR7 catalogue (9999 = not available in SDSS-DR7); (11) DR7 g model magnitude corrected for extinction; (12) DR7 r model magnitude corrected for extinction; (13) DR7 i model magnitude corrected for extinction; (14) DR7 u model magnitude corrected for extinction; (15) DR7 z model magnitude corrected for extinction; (16) rest frame $B-V$ color computed using the MGC-SDSS $g-r$ color; (17) rest frame $B-V$ color computed using DR7 $g-r$ color; (18) absolute magnitude in the B band; (19) absolute magnitude in V band; (20) 2PIGG galaxy serial number (9999 if not present in 2PIGG); (21) 2PIGG galaxy group (0 if it is ungrouped, 9999 if galaxy not present in 2PIGG); (22) 2PIGG match flag of hosting group as in column (12) of Table 2; (23) edge flag of the hosting group as in column (13) of Table 2.

The catalogues for "field-single" and "field-binary" galaxies, see Tables 4 and 5 for a subsample (corrisponding to Table 4 and 5 in electronic version for the total catalogues) include the following columns: (1) galaxy serial number in MGC; (2) galaxy redshift in MGC; (3-4) right ascension and declination at epoch J2000 (in degrees); (5) logarithm of galaxy stellar mass (Kroupa IMF) computed using MGC-SDSS colors; (6) logarithm of galaxy stellar mass (Kroupa IMF) we computed using DR7-CAS colors, (7) logarithm of galaxy stellar mass given in the SDSS-DR7 catalogue (9999 = not available in SDSS-DR7); (8) DR7 g model magnitude corrected for extinction; (9) DR7 r model magnitude corrected for extinction; (10) DR7 i model magnitude corrected for extinction; (11) DR7 u model magnitude corrected for extinction; (12) DR7 z model magnitude corrected for extinction; (13) rest frame $B-V$ color computed using the MGC-SDSS $g-r$ color, (14) rest frame $B-V$ color using the DR7 $g-r$ color; (15) absolute magnitude in B band; (16) absolute magnitude in B band; (17) 2PIGG galaxy serial number (9999 if not present in 2PIGG), (18) 2PIGG host group, or groups, associated (0 if it is ungrouped, 9999 if galaxy not present in 2PIGG).

Finally, the "general-field" catalogue in Table 6 for a subsample (corrisponding to Table 6 in electronic version for the total catalogue) has the same entries as the "field-single" and "field-binary" catalogues, with the addition of a column (19) listing the environment to which the galaxy belongs (the group number in PM2GC=group, 1=field-single, 2=field-binary, 3= group at $z<0.04$ or $z>0.1$, 4= members of trial group, but not in final group).

9 SUMMARY AND CONCLUSIONS

Applying a FoF algorithm to a complete spectroscopic sample of galaxies brighter than M_B=-18.7, in the redshift range $0.03 \leqslant z \leqslant 0.11$, taken from MGC survey, we have built the Padova-




Figure 7. Left. The comparison of redshifts between the 81 groups matching a 2PIGG group. Our errors are computed by propagating the error on the velocities and in many case they are as small as point size. The black line is the 1:1 relation **Right.** The comparison between the velocity dispersions. As the error bars are too large and to avoid visual confusion we plotted in the bottom right part of the panel the median error of our velocity dispersion ($\sim$120 $\mathrm{km\,s^{-1}}$) in y and the error used by 2PIGG (85 $\mathrm{km\,s^{-1}}$) in x. The black line is the 1:1 relation.

Millennium Galaxy and Group Catalogue (PM2GC), a galaxy sample representative of the general field population in the local Universe. In the paper we have first described the FoF group-finding method, calibrated on similar parameters used to identify groups at high redshift, and then we have presented the properties of the group and galaxy catalogues obtained.

1057 galaxies belong to 176 groups containing at least three members at $0.04 \leqslant z \leqslant 0.1$. In addition to the group catalogue containing the main group properties (redshift, velocity dispersion, geometric centre etc.), we provide catalogues for galaxies in groups, galaxies in binary systems and isolated galaxies. The binary and single catalogues contain 490 and 1141 galaxies, respectively, at $0.03 \leqslant z \leqslant 0.11$. We also present a general field (GF) catalogue, which comprises all 3210 galaxies at $0.03 \leqslant z \leqslant 0.11$, representative of the general field population, including 522 galaxies that either are in groups at $z < 0.04$ or $z > 0.1$ or are located in a trial group, but not in the final group sample.

We have also determined the galaxy stellar masses for all galaxies in the PM2GC catalogue both using MGC-SDSS magnitudes and SDSS-DR7 CAS magnitudes, and we have showed the existence of a good agreement with DR7 mass estimates.

In order to validate the effectiveness of our method and the robustness of our results, we tested our groups comparing them with the 2PIGG catalogue, finding a good correspondence with 2PIGG for about half of our groups, and identifying a significant number of groups not present in 2PIGG, thanks to the higher spectroscopic completeness.

The PM2GC provides a valuable database for a different number of studies which will allow us a better understanding of the environmental influence on galaxy properties. In addition to galaxy stellar masses, we have estimated galaxy morphologies, surface brightness parameters, star formation histories and local galaxy densities for all PM2GC galaxies that will be presented in forthcoming papers. In the first upcoming paper (Calvi et al. 2011b), we will analyze the mass functions and the morphological distributions in different environments comparing the results with those obtained for WINGS clusters (Vulcani et al. 2011). Other studies are in progress, i.e. the analysis of the star formation histories as a function of environment from spectral information, the analysis of the galaxy mass-size relation in the local field and of the variation of the galaxy mass function with local density.

ACKNOWLEDGMENTS

We would like to thank the team of the Millennium Galaxy Catalogue for the excellent dataset they have made available to the community. Special thanks to Joechen Liske, who provided us the DVD version of the MGC database and very useful elucidations. We also would like to thank Vincent Eke for help in using the 2PIGG catalogue, Jarle Brinchmann for clarifications about the DR7 masses, Antonio Cava, Andrea Biviano and the whole WINGS team for useful discussions. BV and BMP acknowledge financial support from ASI Contract I/016/07/0.

IDGRPM2	zPM2	sigPM2 $\mathrm{km\,s^{-1}}$	ersigPM2 $\mathrm{km\,s^{-1}}$	Ngal	raPM2 deg(J2000)	decPM2 deg(J2000)	R200 Mpc	IDGR2P	IDGRgal2P	Gal_in	flag_type	flag_bor_R
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
GR1002	0.0769	481	389	6	161.6724	-0.0394	1.148	2963	2963	6	2	1
GR1007	0.0944	196	150	6	151.4264	0.0807	0.462	4921	4921	3	2	1
GR1025	0.05	88	45	5	172.0366	0.1444	0.213	9999	1303	3	4	1
GR1059	0.0782	222	135	6	182.876	0.0627	0.529	3205	3205	5	2	1
GR1065	0.0774	456	239	5	168.8397	-0.1213	1.088	2982	2982	3	2	2
GR1072	0.0819	279	273	5	200.7232	-0.2424	0.664	3592	3592	1	5	2
GR1100	0.0486	177	223	3	191.2142	-0.0579	0.427	1206	1206	1	2	1
GR1123	0.0823	116	85	3	199.8421	0.1619	0.275	3736	3736	2	2	1
GR1135	0.1001	211	88	4	182.0762	0.0777	0.498	5252	5252	1	4	1
GR1181	0.0987	439	68	7	169.137	0.0621	1.036	5240	5240	5	2	1

Table 2. Subsample of 10 galaxy groups with their properties. Columns: (1) PM2GC group serial number; (2) median group redshift; (3) rest frame velocity dispersion; (4) velocity dispersion error; (5) number of galaxies in group; (6-7) geometric centre right ascension and declination in degrees; (8) R_{200} in Mpc, (9) matched 2PIGG group serial number (9999 if no match with 2PIGG); (10) 2PIGG group, or groups, to which galaxies of our group belong; (11) number of galaxies of our group that are members of the 2PIGG matched group (0 indicate no galaxies in 2PIGG or ungrouped galaxies); (12) 2PIGG match flag (1=good match with 2PIGG, both in geometric centre and $\Delta z \leqslant 0.0007$, one to one group correspondance, all of our galaxies in common with 2PIGG are in the matched 2PIGG group; 2= as 1 but at least one of our group galaxies is either ungrouped for 2PIGG, and/or belongs to a different 2PIGG group, and/or is not present in 2PIGG; 3 = no match with 2PIGG, 2PIGG has none or < 50 per cent of our redshifts; 4= no match with 2PIGG, and 2PIGG has at least 50 per cent of our redshifts; 5= one group for 2PIGG, two or more groups for us; 6= no match with 2PIGG, high σ for number of members; 0= group with likely edge problems by comparison with 2PIGG), (13) Edge flag, see §3 1=no edge problem; 2= edge problem.

IDMGC	zMGC	raMGC deg(J2000)	decMGC deg(J2000)	IDGRPM2	d_Mpc Mpc	d_R200 R_{200}	MassPM2 $\log_{10}M^{\star}$	MassCAS $\log_{10}M^{\star}$	MassDR7 $\log_{10}M^{\star}$	g_cor	r_cor	i_cor
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
1093	0.089	150.2561	0.0216	GR1011	1.162	4.834	10.9	10.83	10.97	17.431	16.472	16.014
1241	0.0657	150.8201	-0.0045	GR1013	0.256	3.044	9.98	9.96	10.04	17.863	17.253	16.909
1371	0.0672	150.8934	0.0761	GR1013	0.244	2.831	10.95	10.92	11.0	16.827	15.846	15.373
1418	0.0976	151.0467	-0.164	GR1145	0.184	0.753	9.11	9.08	9999.0	19.281	19.11	18.984
1426	0.1001	151.0606	-0.1724	GR1145	0.086	0.348	9.45	9.42	9999.0	19.396	18.987	18.736
1468	0.0955	151.0964	-0.213	GR1145	0.261	1.104	10.83	10.75	10.69	17.354	16.627	16.248
1565	0.0674	150.7827	0.0452	GR1013	0.352	4.13	11.37	11.34	11.29	15.436	14.584	14.195
1580	0.0661	150.9395	0.03	GR1013	0.372	4.462	10.84	10.81	10.82	16.886	15.989	15.575
1711	0.0942	151.364	0.1374	GR1007	0.529	7.548	10.26	10.23	9999.0	18.459	17.693	17.329
1825	0.0947	151.441	0.0789	GR1007	0.092	1.333	9.5	9.39	9999.0	19.185	18.853	18.586

IDMGC	u_cor	z_cor	BV	BVcas	MabsB	MabsV	ID2P	GR2P	flag_type	flag_bor_R
(1)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)
1093	19.467	15.654	0.883	0.838	-20.34	-21.223	15635	4397	4	2
1241	19.276	16.645	0.55	0.535	-19.258	-19.808	15392	2219	2	1
1371	18.749	15.003	0.951	0.934	-20.217	-21.168	15368	2219	2	1
1418	20.276	18.968	0.053	0.031	-18.891	-18.945	9999	9999	6	2
1426	20.312	18.675	0.325	0.305	-18.744	-19.069	9999	9999	6	2
1468	18.849	16.003	0.664	0.61	-20.968	-21.632	15271	5069	6	2
1565	17.332	13.923	0.812	0.785	-21.796	-22.608	15410	2219	2	1
1580	18.872	15.254	0.857	0.837	-20.294	-21.151	9999	9999	2	1
1711	19.824	17.04	0.676	0.656	-19.498	-20.174	15183	4921	2	1
1825	20.045	18.374	0.294	0.218	-18.985	-19.279	9999	9999	2	1

Table 3. Subsample of galaxies in groups with their properties. Columns: (1) galaxy serial number in MGC; (2) galaxy redshift in MGC; (3-4) right ascension and declination in MGC at epoch J2000 (in degrees); (5) group number in PM2GC; (6) distance from group geometric centre in Mpc; (7) distance from group geometric centre in R_{200} units; (8) logarithm of galaxy stellar mass (Kroupa IMF) obtained from MGC-SDSS colors (PM2GC mass); (9) logarithm of galaxy stellar mass (Kroupa IMF) we computed using DR7-CAS colors; (10) logarithm of galaxy stellar mass (Kroupa IMF) given in the SDSS-DR7 catalogue (9999 = not available in SDSS-DR7); (11) DR7 g model magnitude corrected for extinction; (12) DR7 r model magnitude corrected for extinction; (13) DR7 i model magnitude corrected for extinction; (14) DR7 u model magnitude corrected for extinction; (15) DR7 z model magnitude corrected for extinction; (16) rest frame $B-V$ color computed using the MGC-SDSS $g-r$ color; (17) rest frame $B-V$ color computed using DR7 $g-r$ color; (18) absolute magnitude in the B band; (19) absolute magnitude in V band; (20) 2PIGG galaxy serial number (9999 if not present in 2PIGG); (21) 2PIGG galaxy group (0 if it is ungrouped, 9999 if galaxy not present in 2PIGG); (22) 2PIGG match flag of hosting group as in column (12) of Table 2; (23) edge flag of the hosting group as in column (13) of Table 2.

IDMGC	zMGC	raMGC deg(J2000)	decMGC deg(J2000)	MassPM2 $\log_{10}M^{\star}$	MassCAS $\log_{10}M^{\star}$	MassDR7 $\log_{10}M^{\star}$	g_cor	r_cor	i_cor	u_cor	z_cor	BV	BVcas	MabsB	MabsV	ID2P	2PGROUP
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
19621	0.0681	170.5799	-0.1446	10.15	10.11	10.08	17.573	17.001	16.664	18.783	16.473	0.519	0.491	-19.798	-20.318	23969	2325
19714	0.0689	170.5439	-0.231	10.38	10.37	10.5	17.871	16.994	16.596	19.862	16.246	0.818	0.813	-19.276	-20.095	23995	2325
28996	0.0971	182.1661	0.2117	9.85	9.83	9999.0	18.446	18.016	17.655	19.432	17.538	0.344	0.329	-19.684	-20.028	27452	5048
29003	0.1017	182.1215	0.2215	9.93	9.91	9999.0	19.491	18.741	18.339	20.709	18.077	0.653	0.636	-18.762	-19.415	9999	9999
9645	0.0823	159.3158	0.0804	10.91	10.92	10.92	16.098	15.499	15.103	17.521	14.894	0.519	0.524	-21.689	-22.207	20547	3752
9715	0.0826	159.3208	0.0722	9.94	9.94	9999.0	18.629	18.007	17.624	19.898	17.335	0.553	0.551	-19.15	-19.704	20544	3752
23429	0.1064	175.4133	0.214	10.8	10.8	10.84	17.963	17.03	16.589	19.582	16.261	0.81	0.808	-20.364	-21.174	26027	0
23444	0.1066	175.4055	0.2344	9.67	9.66	9999.0	19.553	19.023	18.693	20.525	18.587	0.455	0.445	-18.832	-19.288	9999	9999
96374	0.0917	161.97	0.216	10.18	10.22	10.07	17.663	17.211	16.913	18.866	16.789	0.325	0.356	-20.559	-20.885	19353	4661
11804	0.0946	161.9251	0.2837	9.36	9.34	9999.0	19.428	19.058	18.793	20.528	18.783	0.271	0.261	-18.706	-18.977	9999	9999

Table 4. Subsample of galaxies in binary systems catalogue with their properties. Columns: (1) galaxy serial number in MGC; (2) galaxy redshift in MGC; (3-4) right ascension and declination in MGC at epoch J2000 (in degrees); (5) logarithm of galaxy stellar mass (Kroupa IMF) obtained from MGC-SDSS colors (PM2GC mass); (6) logarithm of galaxy stellar mass (Kroupa IMF) we computed using DR7-CAS colors; (7) logarithm of galaxy stellar mass (Kroupa IMF) given in the SDSS-DR7 catalogue (9999 = not available in SDSS-DR7); (8) DR7 g model magnitude corrected for extinction; (9) DR7 r model magnitude corrected for extinction; (10) DR7 i model magnitude corrected for extinction; (11) DR7 u model magnitude corrected for extinction; (12) DR7 z model magnitude corrected for extinction; (13) rest frame $B-V$ color computed using the MGC-SDSS $g-r$ color; (14) rest frame $B-V$ color computed using DR7 $g-r$ color; (15) absolute magnitude in the B band; (16) absolute magnitude in V band; (17) 2PIGG galaxy serial number (9999 if not present in 2PIGG); (18) 2PIGG galaxy group (0 if it is ungrouped, 9999 if galaxy not present in 2PIGG).

IDMGC	zMGC	raMGC deg(J2000)	decMGC deg(J2000)	MassPM2 $\log_{10}M^{\star}$	MassCAS $\log_{10}M^{\star}$	MassDR7 $\log_{10}M^{\star}$	g_cor	r_cor	i_cor	u_cor	z_cor	BV	BVcas	MabsB	MabsV	ID2P	2PGROUP
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
61514	0.0844	215.9539	-0.2483	10.57	10.5	10.53	17.277	16.636	16.252	18.52	16.054	0.559	0.511	-20.708	-21.267	38599	0
27280	0.0975	179.8905	-0.0265	10.2	10.2	9999.0	18.757	18.033	17.606	20.054	17.385	0.612	0.607	-19.591	-20.203	27753	0
11383	0.0635	161.5143	0.1471	10.11	10.05	9.92	16.948	16.566	16.294	18.195	16.202	0.364	0.321	-20.256	-20.62	19581	0
66778	0.1079	220.1103	0.0101	9.91	9.88	9999.0	18.963	18.46	18.168	20.029	18.08	0.436	0.414	-19.487	-19.923	39496	5964
52268	0.1076	207.9901	-0.1881	10.47	10.43	9999.0	19.075	18.14	17.61	20.867	17.26	0.838	0.811	-19.431	-20.27	9999	9999
56234	0.1053	211.4459	0.0196	10.13	10.09	9999.0	18.758	18.115	17.788	20.045	17.576	0.543	0.513	-19.668	-20.211	37628	0
16076	0.0813	167.1715	0.1933	10.56	10.44	10.43	17.105	16.574	16.229	18.357	15.989	0.528	0.447	-20.778	-21.307	21884	0
13606	0.108	164.0227	-0.0388	9.95	9.88	9.74	17.877	17.636	17.419	18.692	17.365	0.158	0.112	-20.595	-20.753	22323	6222
57725	0.0804	213.0875	-0.1577	9.8	9.75	9.75	17.831	17.475	17.193	18.914	17.1	0.28	0.244	-19.793	20.073	37292	0
25661	0.0594	178.081	-0.1864	10.21	10.13	10.05	17.165	16.64	16.34	18.405	16.131	0.549	0.495	-19.838	-20.387	25136	0

Table 5. Subsample of galaxies in single catalogue with their properties. Columns: (1) galaxy serial number in MGC; (2) galaxy redshift in MGC; (3-4) right ascension and declination in MGC at epoch J2000 (in degrees); (5) logarithm of galaxy stellar mass (Kroupa IMF) obtained from MGC-SDSS colors (PM2GC mass); (6) logarithm of galaxy stellar mass (Kroupa IMF) we computed using DR7-CAS colors; (7) logarithm of galaxy stellar mass (Kroupa IMF) given in the SDSS-DR7 catalogue (9999 = not available in SDSS-DR7); (8) DR7 g model magnitude corrected for extinction; (9) DR7 r model magnitude corrected for extinction; (10) DR7 i model magnitude corrected for extinction; (11) DR7 u model magnitude corrected for extinction; (12) DR7 z model magnitude corrected for extinction; (13) rest frame $B-V$ color computed using the MGC-SDSS $g-r$ color; (14) rest frame $B-V$ color computed using DR7 $g-r$ color; (15) absolute magnitude in the B band; (16) absolute magnitude in V band; (17) 2PIGG galaxy serial number (9999 if not present in 2PIGG); (18) 2PIGG galaxy group (0 if it is ungrouped, 9999 if galaxy not present in 2PIGG).

IDMGC	zMGC	raMGC deg(J2000)	decMGC deg(J2000)	MassPM2 $\log_{10}M^{\star}$	MassCAS $\log_{10}M^{\star}$	MassDR7 $\log_{10}M^{\star}$	g_cor	r_cor	i_cor	u_cor	z_cor	BV	BVcas	MabsB	MabsV	ID2P	GR2P	flag_env
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)
61514	0.0844	215.9539	-0.2483	10.57	10.5	10.53	17.277	16.636	16.252	18.52	16.054	0.559	0.511	-20.708	-21.267	38599	0	1
27280	0.0975	179.8905	-0.0265	10.2	10.2	9999.0	18.757	18.033	17.606	20.054	17.385	0.612	0.607	-19.591	-20.203	27753	0	1
11383	0.0635	161.5143	0.1471	10.11	10.05	9.92	16.948	16.566	16.294	18.195	16.202	0.364	0.321	-20.256	-20.62	19581	0	1
66778	0.1079	220.1103	0.0101	9.91	9.88	9999.0	18.963	18.46	18.168	20.029	18.08	0.436	0.414	-19.487	-19.923	39496	5964	1
52268	0.1076	207.9901	-0.1881	10.47	10.43	9999.0	19.075	18.14	17.61	20.867	17.26	0.838	0.811	-19.431	-20.27	9999	9999	1
56234	0.1053	211.4459	0.0196	10.13	10.09	9999.0	18.758	18.115	17.788	20.045	17.576	0.543	0.513	-19.668	-20.211	37628	0	1
16076	0.0813	167.1715	0.1933	10.56	10.44	10.43	17.105	16.574	16.229	18.357	15.989	0.528	0.447	-20.778	-21.307	21884	0	1
13606	0.108	164.0227	-0.0388	9.95	9.88	9.74	17.877	17.636	17.419	18.692	17.365	0.158	0.112	-20.595	-20.753	22323	6222	1
57725	0.0804	213.0875	-0.1577	9.8	9.75	9.75	17.831	17.475	17.193	18.914	17.1	0.28	0.244	-19.793	-20.073	37292	0	1
25661	0.0594	178.081	-0.1864	10.21	10.13	10.05	17.165	16.64	16.34	18.405	16.131	0.549	0.495	-19.838	-20.387	25136	0	1
60827	0.0738	215.59	-0.1769	9.7	9.68	9.61	17.98	17.647	17.405	18.969	17.197	0.282	0.265	-19.534	-19.816	38626	0	1
46480	0.0978	201.6882	-0.1698	9.26	9.22	9999.0	19.369	19.127	18.914	20.648	18.85	0.146	0.113	-18.933	-19.079	9999	9999	1
25285	0.0956	177.4551	-0.2612	9.6	9.49	9999.0	19.203	18.842	18.555	20.306	18.557	0.326	0.25	-19.115	-19.441	9999	9999	1

Table 6. Subsample of galaxies in general field catalogue with their properties. Columns: (1) galaxy serial number in MGC; (2) galaxy redshift in MGC; (3-4) right ascension and declination in MGC at epoch J2000 (in degrees); (5) logarithm of galaxy stellar mass (Kroupa IMF) obtained from MGC-SDSS colors (PM2GC mass); (6) logarithm of galaxy stellar mass (Kroupa IMF) we computed using DR7-CAS colors; (7) logarithm of galaxy stellar mass (Kroupa IMF) given in the SDSS-DR7 catalogue (9999 = not available in SDSS-DR7); (8) DR7 g model magnitude corrected for extinction; (9) DR7 r model magnitude corrected for extinction; (10) DR7 i model magnitude corrected for extinction; (11) DR7 u model magnitude corrected for extinction; (12) DR7 z model magnitude corrected for extinction; (13) rest frame $B-V$ color computed using the MGC-SDSS $g-r$ color; (14) rest frame $B-V$ color computed using DR7 $g-r$ color; (15) absolute magnitude in the B band; (16) absolute magnitude in V band; (17) 2PIGG galaxy serial number (9999 if not present in 2PIGG); (18) 2PIGG galaxy group (0 if it is ungrouped, 9999 if galaxy not present in 2PIGG); (19) environment to which the galaxy belongs (the group number in PM2GC=group, 1=field-single, 2=field-binary, 3= group at z$<$0.04 or z$>$0.1, 4= members of trial group, but not in final group).

REFERENCES

Balogh M.L., Bower R.G., 2003, MxAC, 17, 220
Balogh M.L., Morris S.L., Yee H.K.C., Carlberg R.G., Ellingson E.; 1999, ApJ, 527, 54
Balogh M. et al., 2004, MNRAS, 348, 1355
Barnes J.E., Hernquist L., 1996, ApJ, 471, 115
Beers T.C., Flynn K., Gebhardt K., 1990, AJ, 100, 32
Bell E.F., de Jong R.S., 2001, ApJ, 550, 212
Berlind A.A. et al., 2006, ApJ, 167, 1
Berrier J.C., Stewart K.R., Bullock J.S., Purcell C.W., Barton E.J., Wechsler, R.H., 2009, ApJ, 690, 1292
Bertin E., Arnouts S., 1996, A& AS, 117, 39
Blanton M.R., Roweis S., 2007, AJ, 133, 734
Botzler C.S., Snigula J., Bender R., Hopp U., 2004, MNRAS, 349, 425
Butcher H., Oemler A.Jr., 1984, ApJ, 285, 426
Byrd G., Valtonen M., 1990, ApJ, 350, 89
Carlberg R.G., Yee H.K.C., Morris S.L., Lin H., Hall P.B., Patton D.R., Sawicki M., Shepherd C.W., 2001, ApJ, 552, 427
Cimatti A. et al., 2008, A& A, 482, 21
Cox T.J., Jonsson P., Somerville R.S., Primack J.R., Dekel A., 2008, MNRAS, 384, 386
Cross N.J.G. et al., 2004, MNRAS, 349, 576
Dressler A., 1980, ApJ, 236, 351
Dressler A. et al., 1997, ApJ, 490, 577
Driver S.P., Liske J., Cross N.J.G., De Propris R., Allen P.D., 2005, MNRAS, 360, 81
Eke V.R. et al., 2004, MNRAS, 348, 866
Fasano G. et al., 2006, A& A, 445, 805
Finn R.A. et al., 2005, ApJ, 630, 206
Font A.S. et al., 2008, MNRAS, 389, 1619
Fritz J. et al., 2007, A& A, 470, 137
Fritz J. et al., 2011, A& A, 526, 45
Geller M.J., Huchra J.P., 1983, ApJS, 52, 61
Gerke, B.F. et al., 2005, ApJ, 625, 6
Guo Q. et al., 2011, MNRAS, in press
Gunn J.E., Gott J.R., 1972, ApJ, 176, 1
Halliday C. et al., 2004, A& A, 427, 397
Hashimoto Y., Oemler A.Jr. Lin H., Tucker D.L., 1998, ApJ, 499, 589
Henry J.P. et al., 1995, ApJ, 449, 422
Hickson P., Kindl E., Auman J.R., 1989, ApJS, 70, 687
Hopkins A.M., 2004, ApJ, 615, 209
Huchra J.P., Geller M.J., 1982, ApJ, 257, 4
Irwin M., Lewis J., 2001, NewAR, 45, 105
Kang X., Jing Y.P., Mo H.J., Börner G., 2005, ApJ, 631, 21
Kroupa P., 2001, MNRAS, 322, 231
Lewis I. et al., 2002, MNRAS, 334, 673L
Lilly S.J., Le Fevre O., Hammer F., Crampton D., 1996, ApJ, 460L, 1L
Liske J., Lemon D.J., Driver S.P., Cross N.J.G., Couch W.J., 2003, MNRAS, 344, 307
Liu H.B., Hsieh B.C., Ho P.T.P., Lin L., Yan R., 2008, ApJ, 681, 1046L
Madau P., della Valle M., Panagia N., 1998, MNRAS, 297L, 17
Marinoni C., Davis M., Newman J.A., Coil A.L., 2002, ApJ, 580, 122
Martínez H.J., Zandivarez A., Merchán M.E., Domínguez M.J.L., 2002, MNRAS, 337, 1441
McGee S.L. et al., 2008, MNRAS, 387, 1605
Mihos J.C., 2004, Clusters of galaxies: probes of cosmological structure and galaxy evolution, p.278, eds. J.S. Mulchaey, A.Dressler, A. Oemler, Cambridge University press
Miller C.J. et al., 2005, AJ, 130, 968
Milvang-Jensen B. et al., 2008, A& A, 482, 419
Moore B., Frenk C.S., White S.D.M., 1993, MNRAS, 261, 827
Moore B., Katz N., Lake G., Dressler A., Oemler A., 1996, Nature, 379, 613
Mulchaey J.S., Davis D.S., Mushotzky R.F., Burstein D., 2003, ApJS, 145, 39
Murante G. et al., 2007, MNRAS, 377, 2
Poggianti B.M., 1997, A& AS, 1997, 122, 399P
Poggianti B.M. et al., 1999, ApJ, 518, 576P
Poggianti B.M. et al., 2006, ApJ, 642, 188P
Ramella M., Geller M.J., Huchra J.P., 1989, ApJ, 344, 57
Ramella M., Pisani A., Geller M.J., 1997, AJ, 113, 483
Ramella M. et al., 1999, A& A, 342, 1
Salpeter E.E., 1955, ApJ, 121, 161
Somerville R.S. et al., 2008, MNRAS, 391, 481
Springel V., White S.D.M., Tormen G., Kauffmann G., 2001, MNRAS, 328, 726
Toomre A., Toomre J., 1972, ApJ, 178, 623
Tucker D.L. et al., 2000, ApJS, 130, 237
van Breukelen C. et al., 2006, MNRAS, 373, 26
Vulcani B. et al., 2011, MNRAS, 412, 426
Wang L.L.C., Kauffmann G., De Lucia G., 2007, MNRAS, 377, 1419
Wilman D.J., Balogh M.L., Bower R.G., Mulchaey J.S., Oemler A.Jr., Carlberg R.G., 2005, arXiv:astro-ph/0510077
Weinmann S.M., Kauffmann, G., von der Linden A., De Lucia G., 2010, MNRAS, 406, 2249
White S.D.M. et al., 2005, A& A, 444, 365
Yang X., Mo H.J., van den Bosch F.C., Weinmann S.M., Li C., Jing Y.P., 2005, MNRAS, 362, 711
Zabludoff A.I., Mulchaey J.S., 1998, ApJ, 496, 39